

Complete all the following on notebook paper.

\_\_\_\_\_ 1.

Let  $f$  be a continuous function on the closed interval  $[0, 2]$ . If  $2 \leq f(x) \leq 4$ , then the greatest possible value of  $\int_0^2 f(x) dx$  is

- (A) 0            (B) 2            (C) 4            (D) 8            (E) 16

\_\_\_\_\_ 2.

$$\int_0^1 (3x-2)^2 dx =$$

- (A)  $-\frac{7}{3}$             (B)  $-\frac{7}{9}$             (C)  $\frac{1}{9}$             (D) 1            (E) 3

\_\_\_\_\_ 3.

If  $y = 2 \cos\left(\frac{x}{2}\right)$ , then  $\frac{d^2y}{dx^2} =$

- (A)  $-8 \cos\left(\frac{x}{2}\right)$     (B)  $-2 \cos\left(\frac{x}{2}\right)$     (C)  $-\sin\left(\frac{x}{2}\right)$     (D)  $-\cos\left(\frac{x}{2}\right)$     (E)  $-\frac{1}{2} \cos\left(\frac{x}{2}\right)$

\_\_\_\_\_ 4.

$$\int_2^3 \frac{x}{x^2+1} dx =$$

- (A)  $\frac{1}{2} \ln \frac{3}{2}$             (B)  $\frac{1}{2} \ln 2$             (C)  $\ln 2$             (D)  $2 \ln 2$             (E)  $\frac{1}{2} \ln 5$

\_\_\_\_\_ 5.

The area of the region enclosed by the graphs of  $y = x$  and  $y = x^2 - 3x + 3$  is

- (A)  $\frac{2}{3}$             (B) 1            (C)  $\frac{4}{3}$             (D) 2            (E)  $\frac{14}{3}$


6.

Let  $f$  be a polynomial function with degree greater than 2. If  $a \neq b$  and  $f(a) = f(b) = 1$ , which of the following must be true for at least one value of  $x$  between  $a$  and  $b$ ?

- I.  $f(x) = 0$
- II.  $f'(x) = 0$
- III.  $f''(x) = 0$

- (A) None      (B) I only      (C) II only      (D) I and II only      (E) I, II, and III

7.



The graph of  $y = f(x)$  on the closed interval  $[2, 7]$  is shown above. How many points of inflection does this graph have on this interval?


- (A) One      (B) Two      (C) Three      (D) Four      (E) Five

8.

$$\lim_{x \rightarrow \frac{\pi}{4}} \frac{\sin\left(x - \frac{\pi}{4}\right)}{x - \frac{\pi}{4}} \text{ is}$$

- (A) 0      (B)  $\frac{1}{\sqrt{2}}$       (C)  $\frac{\pi}{4}$       (D) 1      (E) nonexistent

9.



If  $f$  is the continuous, strictly increasing function on the interval  $a \leq x \leq b$  as shown above, which of the following must be true?

I.  $\int_a^b f(x) dx < f(b)(b-a)$

II.  $\int_a^b f(x) dx > f(a)(b-a)$

III.  $\int_a^b f(x) dx = f(c)(b-a)$  for some number  $c$  such that  $a < c < b$

- (A) I only      (B) II only      (C) III only      (D) I and III only      (E) I, II, and III


10.

An equation of the line normal to the graph of  $y = x^3 + 3x^2 + 7x - 1$  at the point where  $x = -1$  is

- (A)  $4x + y = -10$       (B)  $x - 4y = 23$       (C)  $4x - y = 2$       (D)  $x + 4y = 25$       (E)  $x + 4y = -25$

11. 2003—AB5B (No Calculator)

Let  $f$  be a function defined on the closed interval  $[0, 7]$ . The graph of  $f$ , consisting of four line segments, is shown above. Let  $g$  be the function given by  $g(x) = \int_2^x f(t) dt$ .



- Find  $g(3)$ ,  $g'(3)$ , and  $g''(3)$ .
- Find the average rate of change of  $g$  on the interval  $0 \leq x \leq 3$ .
- For how many values  $c$ , where  $0 < c < 3$ , is  $g'(c)$  equal to the average rate found in part (b)? Explain your reasoning.
- Find the  $x$ -coordinate of each point of inflection of the graph of  $g$  on the interval  $0 < x < 7$ . Justify your answer.

12. 2003-AB6B (No Calculator)

Let  $f$  be the function satisfying  $f'(x) = x\sqrt{f(x)}$  for all real numbers  $x$ , where  $f(3) = 25$ .

- Find  $f''(3)$ .
- Write an expression for  $y = f(x)$  by solving the differential equation  $\frac{dy}{dx} = x\sqrt{y}$  with the initial condition  $f(3) = 25$ .