

1.

At time $t \geq 0$, a particle moving in the xy -plane has velocity vector given by $v(t) = \langle t^2, 5t \rangle$. What is the acceleration vector of the particle at time $t = 3$?

- (A) $\langle 9, \frac{45}{2} \rangle$ (B) $\langle 6, 5 \rangle$ (C) $\langle 2, 0 \rangle$ (D) $\sqrt{306}$ (E) $\sqrt{61}$

2.

$$\int xe^{x^2} dx =$$

- (A) $\frac{1}{2}e^{x^2} + C$ (B) $e^{x^2} + C$ (C) $xe^{x^2} + C$ (D) $\frac{1}{2}e^{2x} + C$ (E) $e^{2x} + C$

3.

Consider the series $\sum_{n=1}^{\infty} \frac{e^n}{n!}$. If the ratio test is applied to the series, which of the following inequalities results, implying that the series converges?

- (A) $\lim_{n \rightarrow \infty} \frac{e}{n!} < 1$
(B) $\lim_{n \rightarrow \infty} \frac{n!}{e} < 1$
(C) $\lim_{n \rightarrow \infty} \frac{n+1}{e} < 1$
(D) $\lim_{n \rightarrow \infty} \frac{e}{n+1} < 1$
(E) $\lim_{n \rightarrow \infty} \frac{e}{(n+1)!} < 1$

4.

Which of the following gives the length of the path described by the parametric equations $x = \sin(t^3)$ and $y = e^{5t}$ from $t = 0$ to $t = \pi$?

(A) $\int_0^\pi \sqrt{\sin^2(t^3) + e^{10t}} dt$

(B) $\int_0^\pi \sqrt{\cos^2(t^3) + e^{10t}} dt$

(C) $\int_0^\pi \sqrt{9t^4 \cos^2(t^3) + 25e^{10t}} dt$

(D) $\int_0^\pi \sqrt{3t^2 \cos(t^3) + 5e^{5t}} dt$

(E) $\int_0^\pi \sqrt{\cos^2(3t^2) + e^{10t}} dt$

5.

$$f(x) = \begin{cases} \frac{x^2 - 4}{x - 2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$

Let f be the function defined above. Which of the following statements about f are true?

I. f has a limit at $x = 2$.

II. f is continuous at $x = 2$.

III. f is differentiable at $x = 2$.

(A) I only

(B) II only

(C) III only

(D) I and II only

(E) I, II, and III

6.

Given that $y(1) = -3$ and $\frac{dy}{dx} = 2x + y$, what is the approximation for $y(2)$ if Euler's method is used with a step size of 0.5, starting at $x = 1$?

- (A) -5 (B) -4.25 (C) -4 (D) -3.75 (E) -3.5

7.

x	2	3	5	8	13
$f(x)$	6	-2	-1	3	9

The function f is continuous on the closed interval $[2, 13]$ and has values as shown in the table above. Using the intervals $[2, 3]$, $[3, 5]$, $[5, 8]$, and $[8, 13]$, what is the approximation of $\int_2^{13} f(x) dx$ obtained from a left Riemann sum?

- (A) 6 (B) 14 (C) 28 (D) 32 (E) 50

8.

In the xy -plane, what is the slope of the line tangent to the graph of $x^2 + xy + y^2 = 7$ at the point $(2, 1)$?

- (A) $-\frac{4}{3}$ (B) $-\frac{5}{4}$ (C) -1 (D) $-\frac{4}{5}$ (E) $-\frac{3}{4}$

9.

Let R be the region between the graph of $y = e^{-2x}$ and the x -axis for $x \geq 3$. The area of R is

- (A) $\frac{1}{2e^6}$ (B) $\frac{1}{e^6}$ (C) $\frac{2}{e^6}$ (D) $\frac{\pi}{2e^6}$ (E) infinite

10.

Which of the following series converges for all real numbers x ?

- (A) $\sum_{n=1}^{\infty} \frac{x^n}{n}$
- (B) $\sum_{n=1}^{\infty} \frac{x^n}{n^2}$
- (C) $\sum_{n=1}^{\infty} \frac{x^n}{\sqrt{n}}$
- (D) $\sum_{n=1}^{\infty} \frac{e^n x^n}{n!}$
- (E) $\sum_{n=1}^{\infty} \frac{n! x^n}{e^n}$

11. (2013, BC-6)

A function f has derivatives of all orders at $x = 0$. Let $P_n(x)$ denote the n th-degree Taylor polynomial for f about $x = 0$.

- (a) It is known that $f(0) = -4$ and that $P_1\left(\frac{1}{2}\right) = -3$. Show that $f'(0) = 2$.
- (b) It is known that $f''(0) = -\frac{2}{3}$ and $f'''(0) = \frac{1}{3}$. Find $P_3(x)$.
- (c) The function h has first derivative given by $h'(x) = f(2x)$. It is known that $h(0) = 7$. Find the third-degree Taylor polynomial for h about $x = 0$.

12. (2010, BC-6B)

The Maclaurin series for the function f is given by $f(x) = \sum_{n=2}^{\infty} \frac{(-1)^n (2x)^n}{n-1}$ on its interval of convergence.

(a) Find the interval of convergence for the Maclaurin series of f . Justify your answer.

(b) Show that $y = f(x)$ is a solution to the differential equation $xy' - y = \frac{4x^2}{1+2x}$ for $|x| < R$, where R is the radius of convergence from part (a).