

1.

If $f(x) = (\ln x)^2$, then $f''(\sqrt{e}) =$

- (A) $\frac{1}{e}$ (B) $\frac{2}{e}$ (C) $\frac{1}{2\sqrt{e}}$ (D) $\frac{1}{\sqrt{e}}$ (E) $\frac{2}{\sqrt{e}}$

2.

What are all values of x for which the series $\sum_{n=1}^{\infty} \left(\frac{2}{x^2 + 1}\right)^n$ converges?

- (A) $-1 < x < 1$
(B) $x > 1$ only
(C) $x \geq 1$ only
(D) $x < -1$ and $x > 1$ only
(E) $x \leq -1$ and $x \geq 1$

3.

Let h be a differentiable function, and let f be the function defined by $f(x) = h(x^2 - 3)$. Which of the following is equal to $f'(2)$?

- (A) $h'(1)$ (B) $4h'(1)$ (C) $4h'(2)$ (D) $h'(4)$ (E) $4h'(4)$

4.

In the xy -plane, the line $x + y = k$, where k is a constant, is tangent to the graph of $y = x^2 + 3x + 1$. What is the value of k ?

- (A) -3 (B) -2 (C) -1 (D) 0 (E) 1

5.

$$\int \frac{7x}{(2x-3)(x+2)} dx =$$

- (A) $\frac{3}{2} \ln|2x-3| + 2 \ln|x+2| + C$
(B) $3 \ln|2x-3| + 2 \ln|x+2| + C$
(C) $3 \ln|2x-3| - 2 \ln|x+2| + C$
(D) $-\frac{6}{(2x-3)^2} - \frac{2}{(x+2)^2} + C$
(E) $-\frac{3}{(2x-3)^2} - \frac{2}{(x+2)^2} + C$

6.

What is the sum of the series $1 + \ln 2 + \frac{(\ln 2)^2}{2!} + \dots + \frac{(\ln 2)^n}{n!} + \dots$?

(A) $\ln 2$

(B) $\ln(1 + \ln 2)$

(C) 2

(D) e^2

(E) The series diverges.

7.

x	0	1
$f(x)$	2	4
$f'(x)$	6	-3
$g(x)$	-4	3
$g'(x)$	2	-1

The table above gives values of f , f' , g , and g' for selected values of x . If $\int_0^1 f'(x)g(x) dx = 5$, then

$$\int_0^1 f(x)g'(x) dx =$$

(A) -14

(B) -13

(C) -2

(D) 7

(E) 15

8.

If $f(x) = x \sin(2x)$, which of the following is the Taylor series for f about $x = 0$?

(A) $x - \frac{x^3}{2!} + \frac{x^5}{4!} - \frac{x^7}{6!} + \dots$


(B) $x - \frac{4x^3}{2!} + \frac{16x^5}{4!} - \frac{64x^7}{6!} + \dots$

(C) $2x - \frac{8x^3}{3!} + \frac{32x^5}{5!} - \frac{128x^7}{7!} + \dots$

(D) $2x^2 - \frac{2x^4}{3!} + \frac{2x^6}{5!} - \frac{2x^8}{7!} + \dots$

(E) $2x^2 - \frac{8x^4}{3!} + \frac{32x^6}{5!} - \frac{128x^8}{7!} + \dots$

9.



Which of the following differential equations for a population P could model the logistic growth shown in the figure above?

(A) $\frac{dP}{dt} = 0.2P - 0.001P^2$

(B) $\frac{dP}{dt} = 0.1P - 0.001P^2$

(C) $\frac{dP}{dt} = 0.2P^2 - 0.001P$

(D) $\frac{dP}{dt} = 0.1P^2 - 0.001P$

(E) $\frac{dP}{dt} = 0.1P^2 + 0.001P$

10.

In the xy -plane, a particle moves along the parabola $y = x^2 - x$ with a constant speed of $2\sqrt{10}$ units per second.

If $\frac{dx}{dt} > 0$, what is the value of $\frac{dy}{dt}$ when the particle is at the point $(2, 2)$?

- (A) $\frac{2}{3}$ (B) $\frac{2\sqrt{10}}{3}$ (C) 3 (D) 6 (E) $6\sqrt{10}$


11. (2012, BC-6)

The function g has derivatives of all orders, and the Maclaurin series for g is

$$\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+3} = \frac{x}{3} - \frac{x^3}{5} + \frac{x^5}{7} - \dots$$

- (a) Using the ratio test, determine the interval of convergence of the Maclaurin series for g .
- (b) The Maclaurin series for g evaluated at $x = \frac{1}{2}$ is an alternating series whose terms decrease in absolute value to 0. The approximation for $g\left(\frac{1}{2}\right)$ using the first two nonzero terms of this series is $\frac{17}{120}$. Show that this approximation differs from $g\left(\frac{1}{2}\right)$ by less than $\frac{1}{200}$.
- (c) Write the first three nonzero terms and the general term of the Maclaurin series for $g'(x)$.

12. (2011, BC-6)



Let $f(x) = \sin(x^2) + \cos x$. The graph of $y = |f^{(5)}(x)|$ is shown above.

- (a) Write the first four nonzero terms of the Taylor series for $\sin x$ about $x = 0$, and write the first four nonzero terms of the Taylor series for $\sin(x^2)$ about $x = 0$.
- (b) Write the first four nonzero terms of the Taylor series for $\cos x$ about $x = 0$. Use this series and the series for $\sin(x^2)$, found in part (a), to write the first four nonzero terms of the Taylor series for f about $x = 0$.
- (c) Find the value of $f^{(6)}(0)$.
- (d) Let $P_4(x)$ be the fourth-degree Taylor polynomial for f about $x = 0$. Using information from the graph of $y = |f^{(5)}(x)|$ shown above, show that $\left| P_4\left(\frac{1}{4}\right) - f\left(\frac{1}{4}\right) \right| < \frac{1}{3000}$.