

- |      |       |
|------|-------|
| 1. E | 6. E  |
| 2. D | 7. C  |
| 3. D | 8. A  |
| 4. A | 9. E  |
| 5. D | 10. D |

6

1. Let  $f$  and  $g$  be differentiable functions with the following properties:

(i)  $g(x) > 0$  for all  $x$

(ii)  $f(0) = 1$


If  $h(x) = f(x)g(x)$  and  $h'(x) = f(x)g'(x)$ , then  $f'(x) =$

- (A)  $f'(x)$  (B)  $g(x)$  (C)  $e^x$  (D) 0 (E) 1

$h'(x) = f'(x)g(x) + f(x)g'(x) = f(x)g'(x)$

2. The flow of oil, in barrels per hour, through a pipeline on July 9 is given by the graph shown at right. Of the following, which best approximates the total number of barrels of oil that passed through the pipeline that day?

- (A) 500 (B) 600 (C) 2,400 (D) 3,000 (E) 4,800



3. What is the instantaneous rate of change at  $x = 2$  of the function  $f$  given by  $f(x) = \frac{x^2 - 2}{x - 1}$ ?

- (A) -2 (B)  $\frac{1}{6}$  (C)  $\frac{1}{2}$  (D) 2 (E) 6

4. If  $f$  is a linear function and  $0 < a < b$ , then  $\int_a^b f''(x) dx =$

$f(x) = mx + b$   
 $f'(x) = m$

(A) 0 (B) 1 (C)  $\frac{ab}{2}$  (D)  $b - a$  (E)  $\frac{b^2 - a^2}{2}$

$f'(b) - f'(a) = m - m = 0$

5. If  $F(x) = \int_0^x \sqrt{t^3 + 1} dt$ , then  $F'(2) =$


- $F'(x) = \sqrt{x^3 + 1}$  (A) -3 (B) -2 (C) 2 (D) 3 (E) 18

6. If  $f(x) = \sin(e^{-x})$ , then  $f'(x) =$

- (A)  $-\cos(e^{-x})$  (B)  $\cos(e^{-x}) + e^{-x}$  (C)  $\cos(e^{-x}) - e^{-x}$  (D)  $e^{-x} \cos(e^{-x})$  (E)  $-e^{-x} \cos(e^{-x})$

7. If  $f''(x) = x(x+1)(x-2)^2$ , then the graph of  $f$  has inflection points when  $x =$

- (A) -1 only (B) 2 only (C) -1 and 0 only (D) -1 and 2 only (E) -1, 0, and 2 only



8. What are all the values of  $k$  for which  $\int_{-3}^k x^2 dx = 0$ ?

- (A) -3 (B) 0 (C) 3 (D) -3 and 3 (E) -3, 0, and 3


$$\text{Avg} = \frac{\int_1^6 2e^{x-3} dx}{6-1} = \frac{2}{5} e^{x-3} \Big|_1^6 = \frac{2}{5} (e^3 - e^{-2})$$

9. The average value of the function  $f(x) = 2e^{(x-3)}$  on the interval  $[1, 6]$  is

- (A)  $\frac{e^3}{3}$  (B)  $2e^3 - 2e^{-2}$  (C)  $\frac{e^3 - e^2}{3}$  (D)  $e^3 + e^{-5}$  (E)  $\frac{2e^3 - 2e^{-2}}{5}$

10. A rectangle has its base on the  $x$ -axis and both its other vertices on the positive portion of the parabola  $y = 3 - 4x^2$ . What is the maximum possible area of this rectangle?

- (A)  $\frac{3\sqrt{6}}{4}$  (B)  $\frac{3\sqrt{15}}{5}$  (C)  $\frac{3\sqrt{15}}{10}$  (D) 2 (E)  $\frac{3}{2}$



11. (2003-AB2) (Calculator Permitted) A particle moves along the  $x$ -axis so that its velocity at time  $t$  is given

by  $v(t) = -(t+1)\sin\left(\frac{t^2}{2}\right)$ . At time  $t = 0$ , the particle is at position  $x = 1$ .

Handwritten notes for problem 10:

$$y = 3 - 4x^2$$

$$x^2 = \frac{3-y}{4}$$

$$x = \pm \frac{1}{2}\sqrt{3-y}$$


$$A = 2x \cdot y = (1)\sqrt{3-y}$$

$$A = 2x(3-4x^2)$$

$$A = 6x - 8x^3$$

$$A' = 6 - 24x^2 = 0$$

- (a) Find the acceleration of the particle at time  $t = 2$ . Is the speed of the particle increasing at  $t = 2$ ? Why or why not?
- (b) Find all times  $t$  in the open interval  $0 < t < 3$  when the particle changes direction. Justify your answer.
- (c) Find the total distance traveled by the particle from time  $t = 0$  until time  $t = 3$ .
- (d) During the time interval  $0 \leq t \leq 3$ , what is the greatest distance between the particle and the origin? Show the work that leads to your answer.



$t$ (minutes)	$R(t)$ (gallons per minute)
0	20
30	30
40	40
50	55
70	65
90	70

12. (2003-AB3) (Calculator Permitted) The rate of fuel consumption, in gallons per minute, recorded during an airplane flight is given by a twice-differentiable and strictly increasing function  $R$  of time  $t$ . The graph of  $R$  and a table of selected values of  $R(t)$ , for the time interval  $0 \leq t \leq 90$  minutes, are shown above.

- (a) Use data from the table to find an approximation for  $R'(45)$ . Show the computations that lead to your answer. Indicate units of measure.
- (b) The rate of fuel consumption is increasing fastest at time  $t = 45$  minutes. What is the value of  $R''(45)$ ? Explain your reasoning.

- (c) Approximate the value of  $\int_0^{90} R(t)dt$  using a left Riemann sum with five subintervals indicated by the

data in the table. Is this numerical approximation less than the value of  $\int_0^{90} R(t)dt$ ? Explain your reasoning.

- (d) For  $0 < b \leq 90$  minutes, explain the meaning of  $\int_0^b R(t)dt$  in terms of fuel consumption for the plane.

Explain the meaning of  $\frac{1}{b} \int_0^b R(t)dt$  in terms of fuel consumption for the plane. Indicate units of measure in both answers.

