


§8.2—Polar Area

We are going to look at areas enclosed by polar curves, that's *enclosed*, not *under* as we typically have in these problems. These problems work a little differently in polar coordinates. Here is a sketch of what the area that we'll be finding in this section looks like.



The formula for polar area is different from all previous area formulas, because it is not based on rectangles. Instead, polar area uses an infinite number of sectors to find the area. Remember that a sector is a hunk of a circle, a slice of pizza from the whole pizza.

The area of a sector of a circle is given by

$A = \frac{1}{2}\theta r^2$, where θ is in radians. Our area is

bounded by the radial lines from $\theta = \alpha$ to $\theta = \beta$

$$Area = \frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta$$

Example 1:

Find the area bounded by the graph of $r = 2 + 2\sin \theta$.

Example 2:

Find the area of one petal of $r = 2 \sin 3\theta$.

Example 3:

Find the area of one petal of $r = 4 \cos 2\theta$.

Example 4:

Find the area inside one loop of $r^2 = 4 \cos 2\theta$.

Example 5:

Find the area inside $r = 3\sin\theta$ and outside $r = 2 - \sin\theta$.

Example 6:

Find the area of the common interior of $r = 3\cos\theta$ and $r = 1 + \cos\theta$.

Example 7:

(Calculator Permitted) A polar curve is defined by the equation $r = \theta + \sin 2\theta$ for $0 \leq \theta \leq \pi$.

(a) Find the area bounded by the curve and the x -axis.


(b) Find the angle θ that corresponds to the point on the curve where $x = -2$.

(c) For $\frac{\pi}{3} < \theta < \frac{2\pi}{3}$, $\frac{dr}{d\theta}$ is negative. What does this say about the graph on this interval?

(d) At what angle θ in the interval $0 \leq \theta \leq \frac{\pi}{2}$ is the curve farthest away from the origin. Justify your answer.

Example 8:

2013 BC-2 (Calculator Permitted)



14. The graphs of the polar curves $r = 3$ and $r = 4 - 2\sin\theta$ are shown in the figure above. The curves intersect when $\theta = \frac{\rho}{6}$ and $\theta = \frac{5\rho}{6}$.
- (a) Let S be the shaded region that is inside the graph of $r = 3$ and also inside the graph of $r = 4 - 2\sin\theta$. Find the area of S .
- (b) A particle moves along the polar curve $r = 4 - 2\sin\theta$ so that at time t seconds, $\theta = t^2$. Find the time t in the interval $1 \leq t \leq 2$ for which the x -coordinate of the particle's position is -1 .
- (c) For the particle described in part (b), find the position vector in terms of t . Find the velocity vector at time $t = 1.5$.